



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2002

---

## **Differential Inhibition of Constitutive and Inducible Nitric Oxide Synthase in Vascular Endothelial Cells by Analogues of Tetrahydrobiopterin**

Mueller, Nicolas J ; Walter, Roland B ; Linscheid, Philippe ; Schaffner, Andreas ; Schoedon, Gabriele

**Abstract:** In the vasculature, a physiologic production of nitric oxide (NO) is maintained by endothelial nitric oxide synthase (eNOS). Induction of inducible nitric oxide synthase (iNOS) under inflammatory conditions (e.g. septic shock) resulting in high levels of nitric oxide (NO) is believed to be partly responsible for the pathophysiologic changes in the vascular system that occur under inflammatory conditions (e.g. septic shock). Both NOS isoforms are dependent on the obligatory cofactor tetrahydrobiopterin (BH4). We investigated the selectivity and potency of the BH4 analogues 4-amino-BH4 and 5-methyl-BH4 in inhibiting eNOS and iNOS in a murine vascular endothelial cell (MVEC) model expressing either eNOS or iNOS under physiologic and inflammatory conditions, respectively. Exogenous BH4 and its precursor sepiapterin both enhanced physiologic eNOS activity in resting MVEC, while 4-amino-BH4 slightly inhibited eNOS. 5-methyl-BH4 did not have any effect on eNOS. BH4, sepiapterin, and 5-methyl-BH4 had no effect on iNOS in inflammatory activated MVEC. In contrast, 4-amino-BH4 selectively inhibited iNOS with a potency comparable to the unselective NOS inhibitor N-monomethyl-L-arginine (L-NMMA). The present study demonstrates that 4-amino-BH4 selectively and potently inhibits iNOS in vascular endothelial cells, while its effect on eNOS is minimal. The selective inhibition of iNOS is a promising strategy for the treatment of inflammatory conditions with high output of NO. Further in vivo studies are required to determine whether inhibition of NO production by analogues of BH4 offers any advantage compared to inhibition by L-arginine analogues

DOI: <https://doi.org/10.1515/pteridines.2002.13.4.126>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-155364>

Journal Article

Published Version

Originally published at:

Mueller, Nicolas J; Walter, Roland B; Linscheid, Philippe; Schaffner, Andreas; Schoedon, Gabriele (2002). Differential Inhibition of Constitutive and Inducible Nitric Oxide Synthase in Vascular Endothelial Cells by Analogues of Tetrahydrobiopterin. *Pteridines*:n/a.

DOI: <https://doi.org/10.1515/pteridines.2002.13.4.126>

Pteridines

Vol. 13, 2002, pp. 126 - 132

## Differential Inhibition of Constitutive and Inducible Nitric Oxide Synthase in Vascular Endothelial Cells by Analogues of Tetrahydrobiopterin

Nicolas J Mueller<sup>1</sup>, Roland B Walter<sup>2,3</sup>, Philippe Linscheid<sup>2</sup>, Andreas Schaffner<sup>2</sup>, Gabriele Schoedon<sup>2</sup>

<sup>1</sup>Infectious Diseases Division, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

<sup>2</sup>Department of Medicine, Research Unit Medical Clinic B, University Hospital, Zürich, Switzerland

<sup>3</sup>Present address: Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA, USA.

### Abstract

In the vasculature, a physiologic production of nitric oxide (NO) is maintained by endothelial nitric oxide synthase (eNOS). Induction of inducible nitric oxide synthase (iNOS) under inflammatory conditions (e.g. septic shock) resulting in high levels of nitric oxide (NO) is believed to be partly responsible for the pathophysiologic changes in the vascular system that occur under inflammatory conditions (e.g. septic shock). Both NOS isoforms are dependent on the obligatory cofactor tetrahydrobiopterin (BH<sub>4</sub>). We investigated the selectivity and potency of the BH<sub>4</sub> analogues 4-amino-BH<sub>4</sub> and 5-methyl-BH<sub>4</sub> in inhibiting eNOS and iNOS in a murine vascular endothelial cell (MVEC) model expressing either eNOS or iNOS under physiologic and inflammatory conditions, respectively. Exogenous BH<sub>4</sub> and its precursor sepiapterin both enhanced physiologic eNOS activity in resting MVEC, while 4-amino-BH<sub>4</sub> slightly inhibited eNOS. 5-methyl-BH<sub>4</sub> did not have any effect on eNOS. BH<sub>4</sub>, sepiapterin, and 5-methyl-BH<sub>4</sub> had no effect on iNOS in inflammatory activated MVEC. In contrast, 4-amino-BH<sub>4</sub> selectively inhibited iNOS with a potency comparable to the unselective NOS inhibitor N $\omega$ -monomethyl-L-arginine (L-NMMA). The present study demonstrates that 4-amino-BH<sub>4</sub> selectively and potently inhibits iNOS in vascular endothelial cells, while its effect on eNOS is minimal. The selective inhibition of iNOS is a promising strategy for the treatment of inflammatory conditions with high output of NO. Further *in vivo* studies are required to determine whether inhibition of NO production by analogues of BH<sub>4</sub> offers any advantage compared to inhibition by L-arginine analogues.

**Key words:** inflammation; vascular endothelial cell, nitric oxide synthase; pteridine; tetrahydrobiopterin; 4-amino-tetrahydrobiopterin, 5-methyl-tetrahydrobiopterin

### Introduction

Recent years clearly established nitric oxide (NO) as a key mammalian metabolite for signaling in the vascular system in both physiological and pathophysiological states (1). The molecular control of NO synthases (NOS), which catalyze the five electron oxidation of L-arginine in a two-step process yielding L-citrulline and NO, has been elucidated in some detail (2). In the physiologic state, vascular endothelial cells constitutively express eNOS protein, and endothelium-derived NO diffuses into subjacent smooth muscle cells to activate soluble guanylate cyclase thereby increasing cGMP and resulting in vasorelaxation (3).

However, under inflammatory conditions eNOS is downregulated whereas activating stimuli induce the expression of iNOS mRNA and subsequent "high-output" NO production (4). It is of note that the expression of iNOS has been found in murine (5), porcine (6), and rat vascular endothelial cells (7, 8), but not in human vascular endothelial cells (9, 10). In contrast, vascular smooth muscle cells from different species, including rat and human, express iNOS mRNA and functional protein upon stimulation with inflammatory stimuli (11-14).

Tetrahydrobiopterin (BH<sub>4</sub>) is essentially required as cofactor for all NOS isoforms; for a recent review on BH<sub>4</sub>'s functions in the vascular system, see (4). The

Correspondence to: Nicolas J. Mueller MD, Infectious Disease Division, Massachusetts General Hospital, 149-13th Street 5th Floor, Room 5234, Charlestown, MA 02129 USA, Tel. 001 617 724 4997, Fax. 001 617 726 5411, nmuellet@partners.org

exact role of BH<sub>4</sub> in NO catalysis remains intensively debated and may include stabilization of the active conformation of the enzyme, action as an allosteric effector enhancing L-arginine binding, and coupling of L-arginine oxidation to NADPH consumption reducing the ratio of superoxide to NO production, and post-transcriptional stabilization of NO synthase mRNA. Vascular endothelial cells express all enzymes necessary for BH<sub>4</sub> generation. They synthesize and secrete small amounts of BH<sub>4</sub> even in the resting state, and inflammatory stimuli increase BH<sub>4</sub> synthesis manifold in these cells. iNOS in many cell types including vascular smooth muscle cells depends on exogenous BH<sub>4</sub> for full enzymatic activity, rendering NO production susceptible to modulation by exogenous BH<sub>4</sub> or analogues (4). To date, however, it is not clear whether iNOS activity in vascular endothelial cells is also prone to modulation by exogenous BH<sub>4</sub>. Given the high rate of BH<sub>4</sub> synthesis and secretion in these cells, we hypothesized that iNOS in vascular endothelial cells behaves differently compared to iNOS from other cell types devoid of abundant BH<sub>4</sub> synthesis. We therefore investigated the effects of BH<sub>4</sub> and the analogues 4-amino-BH<sub>4</sub> and 5-methyl-BH<sub>4</sub> on NO production in murine vascular endothelial cells expressing both eNOS and iNOS (5).

## Methods

### Chemicals

5,6,7,8-tetrahydro-L-erythro-biopterin (tetrahydro-biopterin, BH<sub>4</sub>), the analogues 4-amino-BH<sub>4</sub> and 5-methyl-BH<sub>4</sub>, and sepiapterin were purchased from Dr. B. Schircks Laboratories (Jona, Switzerland). Murine recombinant interferon- $\gamma$  (IFN- $\gamma$ ) was from Peprotech Inc. (Rocky Hill, NJ, USA). Lipopolysaccharide (LPS, *E. coli* O26:B6, Boivin extraction) was from Difco (Detroit, MI, USA) and Sigma (St. Louis, MO, USA). N<sup>o</sup>-nitro-L-arginine methyl ester (L-NAME) and N<sup>o</sup>-monomethyl-L-arginine (L-NMMA) were obtained from Sigma.

### Cell cultures

Murine vascular endothelial cells (MVEC) (15) were maintained in a humidified 5% CO<sub>2</sub>/95% air atmosphere at 37°C in Iscove's Modified Dulbecco's Medium (Cellgro, Mediatech Inc., Herndon, VA, USA) supplemented with 10% fetal bovine serum (heat inactivated, mycoplasma/virus/endotoxin-tested, Life Technologies, Rockville, MD, USA), L-glutamine (2 mM, Cellgro) and gentamicin (50  $\mu$ g ml<sup>-1</sup>, Sigma). Cells were seeded in 24-well cluster plates (Corning Inc., Corning, NY, USA). Confluent cultures were used for experiments. For inflammatory stimulation,

cells were treated with IFN- $\gamma$  (100 U ml<sup>-1</sup>) and LPS (10  $\mu$ g ml<sup>-1</sup>) for 24 or 48 hours as indicated. Cell viability was assessed by trypan blue exclusion and was 95-98% throughout experiments.

### Determination of NOS isoform mRNA.

Expression of NOS isoforms was determined by reverse-transcriptase PCR using isoform-specific primers as described (16). eNOS, 5'TTCACTCTGCC-CCCAGAGATG3' (sense) and 5'AGAGATCTTCA-CTGCATTGG3' (antisense), yielding a 611 bp product, located at bp 911-1522 of the murine eNOS cDNA sequence (17); iNOS, 5'ATGGCTTGCCCTG-GAAGTTTCT3' (sense) and 5'CCTCTGATGGTGC-CATCGGGC3' (antisense), yielding a 817 bp product, located at bp 192-1009 of the murine iNOS cDNA sequence (18). Both iNOS and eNOS were determined in the same cDNA preparations.

### Measurement of nitric oxide/nitrite

NO production was measured as nitrite (NO<sub>2</sub><sup>-</sup>) in culture supernatants by the Griess method as described (19) with sodium nitrite diluted in complete medium as standard. The ratio of nitrite to nitrate was assumed to remain constant during all treatment regimens used. The detection limit for NO<sub>2</sub><sup>-</sup> was 10 pm/well or 1.0  $\mu$ mol L<sup>-1</sup>.

### Statistical analysis

Results are presented as mean  $\pm$  SD. One-way analysis of variance (ANOVA) for repeated measures with Tukey-Kramer Multiple Comparisons Tests were performed using InStat version 3.00 (GraphPad, San Diego, CA, USA).  $p < 0.05$  was considered significant.

## Results

### Effects on eNOS and iNOS mRNA expression and NO production during inflammatory stimulation of murine vascular endothelial cells (MVEC)

In accordance with previous findings (5), resting MVEC secreted small amounts of NO, detected as nitrite in culture supernatants. After 48 hours, 15.12 $\pm$ 4.99  $\mu$ M (range: 8.17 to 29.83  $\mu$ M) nitrite were produced by unstimulated MVEC ( $n=26$ ; data not shown). As reported in an earlier study, incubation of these cells with IFN- $\gamma$ /LPS resulted in a strong induction of NO synthesis (5). Accordingly, IFN- $\gamma$  and LPS-stimulated MVEC produced 234.01 $\pm$ 166.33  $\mu$ M nitrite after 48 hours ( $n=19$ , data not shown); interexperimental variations were high, as indicated by the wide range from 62.17 to 642.00  $\mu$ M nitrite. As shown in Fig. 1, reverse-transcriptase PCR experiments revealed expression of eNOS, but not of iNOS mRNA under




Figure 1. Differential expression of NOS isoforms in confluent MVEC. Under resting conditions, only endothelial eNOS mRNA was found by RT-PCR (upper panel). 24h after stimulation with IFN- $\gamma$  (100 U ml<sup>-1</sup>) and LPS (10  $\mu$ g ml<sup>-1</sup>), iNOS mRNA was strongly induced, whereas eNOS mRNA disappeared to undetectable levels (lower panel). Therefore, MVEC cells switch from eNOS activity under resting to iNOS activity under inflammatory condition. Both eNOS and iNOS were determined in the same cDNA preparations.

resting conditions. In contrast, analysis of mRNA 24h after the addition of IFN- $\gamma$  and LPS showed strong expression of iNOS mRNA, whereas eNOS mRNA could no longer be detected. These basic experiments clearly confirm that unstimulated MVEC switch from low output NO production by eNOS to high output NO synthesis by iNOS after stimulation with inflammatory stimuli.

#### *Effects of BH<sub>4</sub> and sepiapterin on NO production by MVEC*

Fig. 2 shows the effects of increasing concentrations of the natural cofactor BH<sub>4</sub> and its substituent sepiapterin, which is converted intracellularly to BH<sub>4</sub> by the salvage pathway (4) on NO generation. Both BH<sub>4</sub> (Fig. 2A) and sepiapterin (Fig. 2B) slightly enhanced NO production by eNOS, e.g. in unstimulated MVEC. Nitrite in culture supernatants increased from  $14.10 \pm 5.33$  in controls to  $18.90 \pm 7.93$  for BH<sub>4</sub> at 200  $\mu$ M,  $n=8$ ,  $p=0.001$ , and from  $13.79 \pm 5.12$  to  $17.42 \pm 4.31$  for 200  $\mu$ M sepiapterin,  $n=4$ ,  $p<0.0001$ . In contrast, BH<sub>4</sub> and sepiapterin failed to influence NO production by iNOS in stimulated MVEC (from  $190.90 \pm 179.83$  to  $158.33 \pm 208.97$  for 200  $\mu$ M BH<sub>4</sub>,  $n=4$ ,  $p$  is not significant, and from  $220.33 \pm 185.46$  to  $203.72 \pm 162.61$  for 200  $\mu$ M sepiapterin,  $n=3$ ,  $p$  is not significant).




Figure 2. Effects of the natural cofactor BH<sub>4</sub> and its substituent sepiapterin on NO production by MVEC. After achievement of confluence, MVEC were incubated without stimulus (open symbols) or treated with IFN- $\gamma$  (100 U ml<sup>-1</sup>) and LPS (10  $\mu$ g ml<sup>-1</sup>), and either BH<sub>4</sub> (A) or sepiapterin (B) was added at the indicated concentrations. NO production was measured as nitrite in culture supernatants after 48 hours in triplicate wells and duplicate measurements. Results are presented as % change  $\pm$  SD from corresponding control wells. \* denotes  $p<0.01$  as compared to corresponding wells treated with L-NAME 10  $\mu$ M or 25  $\mu$ M and  $p<0.05$  as compared to 50  $\mu$ M. \*\*  $p<0.01$  compared to corresponding controls.

#### *Effects of the pteridine analogues 4-amino-BH<sub>4</sub> and 5-methyl-BH<sub>4</sub> on NO production by MVEC*

4-amino-BH<sub>4</sub> slightly decreased NO production in unstimulated MVEC (Fig. 3A), as shown by a decrease of nitrite in culture supernatants from  $18.42 \pm 7.84$  in control wells to  $11.54 \pm 4.64$  for wells treated with 100  $\mu$ M 4-amino-BH<sub>4</sub> and  $12.58 \pm 1.11$  for wells treated with 200  $\mu$ M 4-amino-BH<sub>4</sub>,  $n=4$ ,  $p=0.01$ . In contrast to its slight effect on eNOS, 4-amino-BH<sub>4</sub> strongly and dose-dependently inhibited iNOS in stimulated MVEC (decrease from  $167.61 \pm 20.04$  in controls to  $43.56 \pm 3.66$  for 200  $\mu$ M 4-amino-BH<sub>4</sub>,  $n=3$ ,  $p<0.0001$ ). The other






Figure 3. Effects of the pteridine analogues 4-amino-BH<sub>4</sub> and 5-methyl-BH<sub>4</sub> on NO production by MVEC. After confluence was achieved, MVEC were incubated without stimulus (open symbols) or treated with IFN- $\gamma$  (100 U ml<sup>-1</sup>) and LPS (10  $\mu$ g ml<sup>-1</sup>) and either 4-amino-BH<sub>4</sub> (A) or 5-methyl-BH<sub>4</sub> (B) was added as indicated. NO production was measured as nitrite in culture supernatants after 48 hours in triplicate wells and duplicate measurements. Results are presented as % change  $\pm$  SD from corresponding control wells. \*  $p < 0.05$  compared to corresponding controls; \*\*  $p < 0.01$  compared to corresponding controls; \*\*\*  $p > 0.001$  compared to corresponding controls.

pteridine analogue 5-methyl-BH<sub>4</sub> failed to influence NO production both in unstimulated as well as in stimulated MVEC (Fig. 3B). In experiments with unstimulated cells, the amount of nitrite in supernatants was  $14.89 \pm 3.43$  in control wells and  $13.39 \pm 7.42$  in wells incubated with 200  $\mu$ M 5-methyl-BH<sub>4</sub>,  $n=3$ ,  $p$  is not significant. In stimulated MVEC, supernatant nitrite was  $267.39 \pm 229.91$  in controls and  $238.00 \pm 197.15$  at 200  $\mu$ M 5-methyl-BH<sub>4</sub>,  $n=3$ ,  $p$  is not significant.

#### *Effects of the L-arginine analogues L-NAME and L-NMMA on NO production by MVEC*

Both L-arginine analogues tested, L-NAME and L-NMMA, significantly inhibited NO production in

unstimulated and stimulated MVEC. L-NAME (Fig. 4A) decreased NO generation by eNOS from  $14.22 \pm 4.83$  to  $8.67 \pm 7.04$  at a concentration of 200  $\mu$ M L-NAME,  $n=3$ ,  $p=0.001$ , and by iNOS from  $344.06 \pm 270.57$  to  $188.50 \pm 176.46$  for 200  $\mu$ M L-NAME,  $n=3$ ,  $p=0.001$ . L-NMMA (Fig. 4B) inhibited NO production by eNOS, as shown by a decrease of nitrite in culture supernatants from  $16.04 \pm 2.87$  to  $10.33 \pm 4.53$  for 200  $\mu$ M L-NMMA,  $n=4$ ,  $p=0.01$ , and it decreased NO generation by iNOS from  $228.17 \pm 95.56$  to  $71.17 \pm 18.77$  for 200  $\mu$ M L-NMMA,  $n=3$ ,  $p=0.0005$ .

#### Discussion

This study shows that highly upregulated NO production in MVEC is regulated on the NOS isoform




Figure 4. Effects of the L-arginine analogues L-NAME and L-NMMA on NO production by MVEC. Confluent MVEC were incubated without stimulus (open symbols) or treated with IFN- $\gamma$  (100 U ml<sup>-1</sup>) and LPS (10  $\mu$ g ml<sup>-1</sup>), and increasing concentrations of either L-NAME (A) or L-NMMA (B) were as indicated. NO production was measured as nitrite in culture supernatants after 48 hours in triplicate wells and duplicate measurements. Results are presented as % change  $\pm$  SD from corresponding control wells. \*  $p < 0.05$  compared to corresponding controls; \*\*  $p < 0.01$  compared to corresponding controls; \*\*\*  $p < 0.001$  compared to corresponding controls.

gene transcription level, confirming the "switch model" of NOS isoforms under inflammatory conditions. Using this unique cellular model, we demonstrate the selective inhibition of iNOS by 4-amino-BH<sub>4</sub>, and an independence of vascular endothelial iNOS from exogenous BH<sub>4</sub> for full enzymatic activity.

The data outlined herein unequivocally show that basal NO production of MVEC can solely be attributed to eNOS activity. However, upon stimulation with proinflammatory molecules, iNOS expression is induced while eNOS is downregulated. The reduction of eNOS mRNA and protein levels following activation by cytokines or endotoxin has also been demonstrated in cultured human and bovine endothelial cells, and this effect could at least partially be attributed to enhanced degradation rates of eNOS transcripts (5, 9, 20-23). Therefore, MVEC switch their active NOS isoform in dependence of the presence of inflammatory signals, i.e. from low-output NO production by eNOS under physiological (non-inflammatory) conditions to high-output NO generation by iNOS under inflammatory situations, referred to as "switch model" previously (4). Although eNOS mRNA is downregulated to undetectable levels within several hours after inflammatory stimulation, preexisting eNOS protein may continue to contribute to the total production of NO during the first days under inflammatory conditions (5); however, this seems to be negligible in view of the large amounts of NO generated simultaneously by iNOS.

Resting vascular endothelial cells constitutively synthesize small amounts of BH<sub>4</sub>. Although the rate of enzyme activity and resulting BH<sub>4</sub> synthesis may depend on experimental conditions and probably also on the type of endothelial cell assessed, this basic characteristic has been found both in freshly isolated cells as well as in cells after prolonged culturing (4). In addition, our findings that sepiapterin slightly increased (and additional BH<sub>4</sub> tended to increase) the amount of NO production of vascular endothelial eNOS are in line with observations made by others and underline the regulatory capacity of the pteridine cofactor for endothelial NO production under non-inflammatory conditions. However, *in vivo* data suggest that under physiologic conditions levels endogenous BH<sub>4</sub> are nearly saturating and barely a limiting factor for optimal or near optimal endothelial eNOS activity, and only very high concentrations of BH<sub>4</sub> resulted in endothelium-dependent vasorelaxation in animal and human studies (24, 25).

It is generally believed that iNOS activity largely depends on exogenous BH<sub>4</sub>, i.e. that endogenous BH<sub>4</sub> biosynthesis does not suffice to support iNOS for full enzymatic activity. Although BH<sub>4</sub> biosynthesis is

induced by inflammatory stimuli in cultured cells, intrinsic BH<sub>4</sub> availability has nevertheless been demonstrated to be a limiting factor for iNOS activity in many cell types including macrophages, mesangial cells, vascular smooth muscle, and fibroblasts, rendering NO production susceptible to modulation of intracellular BH<sub>4</sub> levels by inhibition of BH<sub>4</sub> biosynthesis or exogenous BH<sub>4</sub> addition (4). However, the present study shows strikingly different characteristics for vascular endothelial cells. The vascular endothelium produces and secretes abundant amounts of BH<sub>4</sub> upon inflammatory stimulation, as shown for cells of murine (26) and human (27) origin. The increase in *de novo* BH<sub>4</sub> biosynthesis occurs early after inflammatory stimulation and may even lead to transiently increased NO production by eNOS despite the concomitant decrease of total eNOS protein. In view of this high level BH<sub>4</sub> biosynthesis in endothelial cells under inflammatory conditions, it is not surprising that endogenous BH<sub>4</sub> levels are saturating to fully serve endothelial iNOS protein. Moreover, since most of the newly synthesized BH<sub>4</sub> is secreted, probably even vectorially into the basal direction (4), endothelial BH<sub>4</sub> may serve underlying smooth muscle cells with additional cofactor needed for full iNOS activity, thereby accounting for an inflammatory endothelium-derived relaxing factor. On the other hand, BH<sub>4</sub> is a potent antioxidative and reactive oxygen scavenging agent, preventing peroxynitrite formation when present in saturating amounts (28). The high amount of BH<sub>4</sub> produced by endothelial cells in inflammation or sepsis could therefore be beneficial. Thus, an inhibition of iNOS and not of BH<sub>4</sub> synthesis seems desirable.

The 4-amino analogue of BH<sub>4</sub> binds with high affinity and the same potency to both recombinant eNOS and iNOS and strongly interferes with stimulation of the enzymes by exogenous BH<sub>4</sub>, although 4-amino-BH<sub>4</sub> shares the same allosteric effects, namely stabilization of the active dimers and low-to-high spin transition of the heme, as the natural cofactor (29, 30). In contrast to these findings obtained with purified proteins, 4-amino-BH<sub>4</sub> was much less effective in inhibiting NO formation in cultured porcine endothelial cells (expressing eNOS) compared to cultured murine fibroblasts or human adenocarcinoma cells (expressing iNOS) (29). Despite the fact that the effect of 4-amino-BH<sub>4</sub> critically depended on intracellular BH<sub>4</sub> levels, the selectivity towards inhibition of iNOS could not simply be explained by different intracellular BH<sub>4</sub> levels, and therefore, it was suggested that the high-affinity binding of the inhibitor during protein expression may be responsible for the difference, since the turnover of iNOS protein is much more rapid than that of eNOS (half-life approximately 3 hours vs. 20 hours)



(29). Likewise, 4-amino- BH<sub>4</sub> was shown to be fairly selective in inhibiting iNOS in an in vivo vessel strip model of endotoxemia; in this study, the pteridine antagonist had to be added shortly after the artery strips were treated with inflammatory stimuli (31), again suggesting that the apparent selectivity of 4-amino- BH<sub>4</sub> may be related to high-affinity binding of the inhibitor during protein expression.

The present data are in full agreement with those obtained by (29) and (31) and show that a preferential inhibition of the iNOS protein by 4-amino-BH<sub>4</sub> can also be demonstrated in cells expressing both NOS isoforms, thereby excluding the possibility of cell type differences as an explanation for the observed effect. Nevertheless, different situations will be encountered in vivo in the human vasculature, where endothelial BH<sub>4</sub> might compete with exogenous 4-amino-BH<sub>4</sub> for binding on the pteridine-sites of smooth muscle cell iNOS, and in vivo experiments will ultimately be required to determine whether 4-amino-BH<sub>4</sub> displays a "real" selectivity for iNOS thereby offering an interesting therapeutic tool for the treatment of NO overproduction by iNOS under inflammatory conditions in humans.

The 5-methyl-analogue of BH<sub>4</sub> has previously been shown to substitute BH<sub>4</sub> as a cofactor (32); however, at high concentrations of purified NOS, the autoxidation resistant 5-methyl-BH<sub>4</sub> may result in fully coupled NO generation and, subsequently, NO-induced autoinhibition of NOS (33). Therefore, the effect of 5-methyl-BH<sub>4</sub> on net amounts of NO may be critically dependent on the experimental conditions. In the present study we did not find any significant effect of 5-methyl-BH<sub>4</sub> neither on eNOS nor on iNOS activity, rather excluding the use of this analogue to pharmacologically modulate NO production in the vasculature.

In summary, vascular endothelial cells switch from eNOS under resting to iNOS under inflammatory conditions; the concomitant increase in endogenous BH<sub>4</sub> biosynthesis make these cells independent from exogenous sources for the pteridine cofactor for full enzymatic activity. 4-amino-BH<sub>4</sub> was revealed as a preferential inhibitor for iNOS in vascular endothelial cells, whereas 5-methyl-BH<sub>4</sub> was largely ineffective. Further in vivo experiments are required to confirm these observations and to determine whether inhibition of NO production by analogues of BH<sub>4</sub> offers any advantage over inhibition by L-arginine-analogues in the treatment of human inflammatory diseases.

## References

- 1 Cines DB, Poilak ES, Buck CA, Loscalzo J, Zimmerman GA, McEver RP, Pober JS, Wick TM, Konkle BA, Schwartz BS, Barnathan ES, McCrae KR, Hug BA, Schmidt AM, Stern DM: Endothelial cells in physiology and in the pathophysiology of vascular disorders. *Blood* 1998;91:3527-61.
- 2 Papapetropoulos A, Rudic RD, Sessa WC: Molecular control of nitric oxide synthases in the cardiovascular system. *Cardiovasc Res* 1999; 43:509-20.
- 3 Liaudet L, Soriano FG, Szabo C: Biology of nitric oxide signaling. *Crit Care Med* 2000;28:N37-52.
- 4 Walter R, Schaffner A, Schoedon G: Tetrahydrobiopterin in the vascular system. *Pteridines* 2001, 12:93-120.
- 5 Walter R, Schaffner A, Schoedon G: Differential regulation of constitutive and inducible nitric oxide production by inflammatory stimuli in murine endothelial cells. *Biochem Biophys Res Commun* 1994;202:450-5.
- 6 Radomski MW, Palmer RM, Moncada S: Glucocorticoids inhibit the expression of an inducible, but not the constitutive, nitric oxide synthase in vascular endothelial cells. *Proc Natl Acad Sci USA* 1990;87:10043-7.
- 7 Marumo T, Nakaki T, Adachi H, Esumi H, Suzuki H, Saruta T, Kato R: Nitric oxide synthase mRNA in endothelial cells: synergistic induction by interferon- $\gamma$ , tumor necrosis factor- $\alpha$  and lipopolysaccharide and inhibition by dexamethasone. *Jpn J Pharmacol* 1993;63:327-34.
- 8 Kanno K, Hirata Y, Imai T, Iwashina M, Marumo F: Regulation of inducible nitric oxide synthase gene by interleukin-1 beta in rat vascular endothelial cells. *Am J Physiol* 1994;267:H2318-24.
- 9 Rosenkranz-Weiss P, Sessa WC, Milstien S, Kaufman S, Watson CA, Pober JS: Regulation of nitric oxide synthesis by proinflammatory cytokines in human umbilical vein endothelial cells. Elevations in tetrahydrobiopterin levels enhance endothelial nitric oxide synthase specific activity. *J Clin Invest* 1994;93:2236-43.
- 10 Cepinskas G, Lush CW, Kvietys PR: Anoxia/reoxygenation-induced tolerance with respect to polymorphonuclear leukocyte adhesion to cultured endothelial cells. A nuclear factor-kappaB-mediated phenomenon. *Circ Res* 1999;84:103-12.
- 11 Kanno K, Hirata Y, Imai T, Marumo F: Induction of nitric oxide synthase gene by interleukin in vascular smooth muscle cells. *Hypertension* 1993;22:34-9.
- 12 Nunokawa Y, Ishida N, Tanaka S: Cloning of inducible nitric oxide synthase in rat vascular smooth muscle cells. *Biochem Biophys Res Commun* 1993;191:89-94.



- 13 Thorin-Trescases N, Hamilton CA, Reid JL, McPherson KL, Jardine E, Berg G, Bohr D, Dominiczak AF: Inducible L-arginine/nitric oxide pathway in human internal mammary artery and saphenous vein. *Am J Physiol* 1995;268:H1122-32.
- 14 Chester AH, Borland JA, Buttery LD, Mitchell JA, Cunningham DA, Hafizi S, Hoare GS, Springall DR, Polak JM, Yacoub MH: Induction of nitric oxide synthase in human vascular smooth muscle: interactions between proinflammatory cytokines. *Cardiovasc Res* 1998;38:814-21.
- 15 Williams RL, Risau W, Zerwes HG, Drexler H, Aguzzi A, Wagner EF: Endothelioma cells expressing the polyoma middle T oncogene induce hemangiomas by host cell recruitment. *Cell* 1989;57:1053-63.
- 16 Linscheid P, Schaffner A, Schoedon G: Modulation of inducible nitric oxide synthase mRNA stability by tetrahydrobiopterin in vascular smooth muscle cells. *Biochem Biophys Res Commun* 1998;243:137-41.
- 17 Gnanapandithen K, Chen Z, Kau CL, Gorczynski RM, Marsden PA: Cloning and characterization of murine endothelial constitutive nitric oxide synthase. *Biochem Biophys Acta* 1996;1308:103-6.
- 18 Lowenstein CJ, Glatt CS, Bredt DS, Snyder SH: Cloned and expressed macrophage nitric oxide synthase contrasts with the brain enzyme. *Proc Natl Acad Sci USA* 1992;89:6711-5.
- 19 Schoedon G, Schneemann M, Hofer S, Guerrero L, Blau N, Schaffner A: Regulation of the L-arginine-dependent and tetrahydrobiopterin-dependent biosynthesis of nitric oxide in murine macrophages. *Eur J Biochem* 1993;213:833-9.
- 20 Marsden PA, Schappert KT, Chen HS, Flowers M, Sundell CL, Wilcox JN, Lamas S, Michel T: Molecular cloning and characterization of human endothelial nitric oxide synthase. *FEBS Lett* 1992;307:287-93.
- 21 Myers PR, Wright TF, Tanner MA, Adams HR: EDRF and nitric oxide production in cultured endothelial cells: direct inhibition by *E. coli* endotoxin. *Am J Physiol* 1992;262:H710-8.
- 22 MacNaul KL, Hutchinson NI: Differential expression of iNOS and cNOS mRNA in human vascular smooth muscle cells and endothelial cells under normal and inflammatory conditions. *Biochem Biophys Res Commun* 1993;196:1330-4.
- 23 Linscheid P, Schaffner A, Blau N, Schoedon G: Regulation of 6-pyruvoyltetrahydropterin synthase activity and messenger RNA abundance in human vascular endothelial cells. *Circulation* 1998;98:1703-6.
- 24 Heitzer T, Krohn K, Albers S, Meinertz T: Tetrahydrobiopterin improves endothelium-dependent vasodilation by increasing nitric oxide activity in patients with Type II diabetes mellitus. *Diabetologia* 2000;43:1435-8.
- 25 Heitzer T, Brockhoff C, Mayer B, Warnholtz A, Mollnau H, Henne S, Meinertz T, Munzel T: Tetrahydrobiopterin improves endothelium-dependent vasodilation in chronic smokers: evidence for a dysfunctional nitric oxide synthase. *Circ Res* 2000;86:E36-41.
- 26 Walter R, Schaffner A, Blau N, Kierat L, Schoedon G: Tetrahydrobiopterin is a secretory product of murine vascular endothelial cells. *Biochem Biophys Res Commun* 1994;203:1522-6.
- 27 Schoedon G, Schneemann M, Blau N, Edgell CJ, Schaffner A: Modulation of human endothelial cell tetrahydrobiopterin synthesis by activating and deactivating cytokines: new perspectives on endothelium-derived relaxing factor. *Biochem Biophys Res Commun* 1993;196:1343-8.
- 28 Shimizu S, Ishii M, Kawakami Y, Momose K, Yamamoto T: Protective effects of tetrahydrobiopterin against nitric oxide-induced endothelial cell death. *Life Sci* 1998;63:1585-92.
- 29 Schmidt K, Werner-felmayer G, Mayer B, Werner ER: Preferential inhibition of inducible nitric oxide synthase in intact cells by the 4-amino analogue of tetrahydrobiopterin. *Eur J Biochem* 1999;259:25-31.
- 30 Mayer B, Wu C, Gorren AC, Pfeiffer S, Schmidt K, Clark P, Stuehr DJ, Werner ER: Tetrahydrobiopterin binding to macrophage inducible nitric oxide synthase: heme spin shift and dimer stabilization by the potent pterin antagonist 4-amino-tetrahydrobiopterin. *Biochemistry* 1997;36:8422-7.
- 31 Gibrail HD, Dittrich P, Saleh S, Mayer B: Inhibition of endotoxin-induced vascular hyporeactivity by 4-amino-tetrahydrobiopterin. *Br J Pharmacol* 2000;131:1757-65.
- 32 Riethmuller C, Gorren AC, Pitters E, Hemmens B, Habisch HJ, Heales SJ, Schmidt K, Werner ER, Mayer B: Activation of neuronal nitric-oxide synthase by the 5-methyl analog of tetrahydrobiopterin. Functional evidence against reductive oxygen activation by the pterin cofactor. *J Biol Chem* 1999;274:16047-51.
- 33 Gorren AC, Schrammel A, Riethmuller C, Schmidt K, Koesling D, Werner ER, Mayer B: Nitric oxide-induced autoinhibition of neuronal nitric oxide synthase in the presence of the autooxidation-resistant pteridine 5-methyltetrahydrobiopterin. *Biochem J* 2000;347:475-84.